

The Stability of Matter and Quantum Electrodynamics

Elliott H. Lieb

September 2, 2002

1 Foreword

Heisenberg was undoubtedly one of the most important physicists of the 20th century, especially concerning the creation of quantum mechanics. It was, therefore, a great honor and privilege for me to be asked to speak at this symposium since quantum mechanics is central to my own interests and forms the basis of my talk, which is about the quantum theory of matter in the large and its interaction with the quantized radiation field discovered earlier by Planck.

My enthusiastic participation in the scientific part of this symposium was tempered by other concerns, however. Heisenberg has become, by virtue of his importance in the German and world scientific community, an example of the fact that a brilliant scientific and highly cultured mind could coexist with a certain insensitivity to political matters and the way they affected life for his fellow citizens and others. Many opinions have been expressed about his participation in the struggle of the Third Reich for domination, some forgiving and some not, and I cannot judge these since I never met the man. But everyone is agreed about the fact that Heisenberg could view with equanimity, if not some enthusiasm, the possibility of a German victory,

For publication in the proceedings of the Werner Heisenberg Centennial, Munich, December, 2001.

© 2002 by the author. This article may be reproduced, in its entirety, for non-commercial purposes.

Work partially supported by U.S. National Science Foundation grant PHY 0139984.

which clearly would have meant the end of civilization as we know it and enjoy it. By the start of the war this fact was crystal clear, or should have been clear if humanistic culture has more than a superficial meaning. To me it continues to be a mystery that the same person could see the heights of human culture and simultaneously glimpse into the depths of depravity and not see that the latter would destroy the former were it not itself destroyed.

2 Introduction

The quantum mechanical revolution brought with it many successes but also a few problems that have yet to be resolved. We begin with a sketch of the topics that will concern us here.

2.1 Triumph of Quantum Mechanics

One of the basic problems of classical physics (after the discovery of the point electron by Thompson and of the (essentially) point nucleus by Rutherford) was the stability of atoms. Why do the electrons in an atom not fall into the nucleus? Quantum mechanics explained this fact. It starts with the classical Hamiltonian of the system (nonrelativistic kinetic energy for the electrons plus Coulomb's law of electrostatic energy among the charged particles). By virtue of the non-commutativity of the kinetic and potential energies in quantum mechanics the stability of an atom – in the sense of a finite lower bound to the energy – was a consequence of the fact that any attempt to make the electrostatic energy very negative would require the localization of an electron close to the nucleus and this, in turn, would result in an even greater, positive, kinetic energy.

Thus, the basic stability problem for an atom was solved by an inequality that says that $\langle 1/|x| \rangle$ can be made large only at the expense of making $\langle p^2 \rangle$ even larger. In elementary presentations of the subject it is often said that the mathematical inequality that ensures this fact is the famous uncertainty principle of Heisenberg (proved by Weyl), which states that $\langle p^2 \rangle \langle x^2 \rangle \geq (9/8)\hbar^2$ with $\hbar = h/2\pi$ and h = Planck's constant.

While this principle is mathematically rigorous it is actually insufficient for the purpose, as explained, e.g., in [17, 19], and thus gives only a heuristic explanation of the power of quantum mechanics to prevent collapse. A more powerful inequality, such as Sobolev's inequality (9), is needed (see,

e.g., [21]). The utility of the latter is made possible by Schrödinger's representation of quantum mechanics (which earlier was a somewhat abstract theory of operators on a Hilbert space) as a theory of differential operators on the space of square integrable functions on $\mathbb{R}^3$. The importance of Schrödinger's representation is sometimes underestimated by formalists, but it is of crucial importance because it permits the use of functional analytic methods, especially inequalities such as Sobolev's, which are not easily visible on the Hilbert space level. These methods are essential for the developments reported here.

To summarize, the understanding of the stability of atoms and ordinary matter requires a formulation of quantum mechanics with two ingredients:

- A Hamiltonian formulation in order to have a clear notion of a lowest possible energy. Lagrangian formulations, while popular, do not always lend themselves to the identification of that quintessential quantum mechanical notion of a ground state energy.
- A formulation in terms of concrete function spaces instead of abstract Hilbert spaces so that the power of mathematical analysis can be fully exploited.

2.2 Some Basic Definitions

As usual, we shall denote the lowest energy (eigenvalue) of a quantum mechanical system by E_0 . (More generally, E_0 denotes the infimum of the spectrum of the Hamiltonian H in case this infimum is not an eigenvalue of H or is $-\infty$.) Our intention is to investigate arbitrarily large systems, not just atoms. In general we suppose that the system is composed of N electrons and K nuclei of various kinds. Of course we could include other kinds of particles but N and K will suffice here. $N = 1$ for a hydrogen atom and $N = 10^{23}$ for a mole of hydrogen. We shall use the following terminology for two notions of stability:

$$E_0 > -\infty \quad \text{Stability of the first kind,} \quad (1)$$

$$E_0 > C(N + K) \quad \text{Stability of the second kind} \quad (2)$$

for some constant $C \leq 0$ that is independent of N and K , but which may depend on the physical parameters of the system (such as the electron charge

and mass). Usually, $C < 0$, which means that there is a positive binding energy per particle.

Stability of the second kind is absolutely essential if quantum mechanics is going to reproduce some of the basic features of the ordinary material world: The energy of ordinary matter is extensive, the thermodynamic limit exists and the laws of thermodynamics hold. Bringing two stones together might produce a spark, but not an explosion with a release of energy comparable to the energy in each stone. Stability of the second kind does not guarantee the existence of the thermodynamic limit for the free energy, but it is an essential ingredient [20] [17, Sect. V].

It turns out that stability of the second kind cannot be taken for granted, as Dyson discovered [8]. If Coulomb forces are involved, then *the Pauli exclusion principle is essential*. Charged bosons are *not stable* because for them $E_0 \sim -N^{7/5}$ (nonrelativistically) and $E_0 = -\infty$ for large, but finite N (relativistically, see Sect. 4.2).

2.3 The Electromagnetic Field

A second big problem handed down from classical physics was the ‘electromagnetic mass’ of the electron. This poor creature has to drag around an infinite amount of electromagnetic energy that Maxwell burdened it with. Moreover, the electromagnetic field itself is quantized – indeed, that fact alone started the whole revolution.

While quantum mechanics accounted for stability with Coulomb forces and Schrödinger led us to think seriously about the ‘wave function of the universe’, physicists shied away from talking about the wave function of the particles in the universe *and* the electromagnetic field in the universe. It is noteworthy that physicists are happy to discuss the quantum mechanical many-body problem with external electromagnetic fields non-perturbatively, but this is rarely done with the quantized field. The quantized field cannot be avoided because it is needed for a correct description of atomic radiation, the laser, etc. However, the interaction of matter with the quantized field is almost always treated perturbatively or else in the context of highly simplified models (e.g., with two-level atoms for lasers).

The quantized electromagnetic field greatly complicates the stability of matter question. It requires, ultimately, mass and charge renormalizations. At present such a complete theory does not exist, but a theory *must* exist because matter exists and because we have strong experimental evidence

about the manner in which the electromagnetic field interacts with matter, i.e., we know the essential features of a low energy Hamiltonian. In short, nature tells us that it must be possible to formulate a self-consistent quantum electrodynamics (QED) *non-perturbatively*, (perhaps with an ultraviolet cutoff of the field at a few MeV). It should not be necessary to have recourse to quantum chromodynamics (QCD) or some other high energy theory to explain ordinary matter.

Physics and other natural sciences are successful because physical phenomena associated with each range of energy and other parameters are explainable to a good, if not perfect, accuracy by an appropriate self-consistent theory. This is true whether it be hydrodynamics, celestial dynamics, statistical mechanics, etc. If low energy physics (atomic and condensed matter physics) is not explainable by a self-consistent, non-perturbative theory on its own level one can speak of an epistemological crisis.

Some readers might say that QED is in good shape. After all, it accurately predicts the outcome of some very high precision experiments (Lamb shift, g -factor of the electron). But the theory does not really work well when faced with the problem, which is explored here, of understanding the many-body ($N \approx 10^{23}$) problem and the stable low energy world in which we spend our everyday lives.

2.4 Relativistic Mechanics

When the classical kinetic energy $p^2/2m$ is replaced by its relativistic version $\sqrt{p^2c^2 + m^2c^4}$ the stability question becomes much more complicated, as will be seen later. It turns out that even stability of the first kind is not easy to obtain and it depends on the values of the physical constants, notably the fine structure constant

$$\alpha = e^2/\hbar c = 1/137.04 , \quad (3)$$

where $-e$ is the electric charge of the electron.

For ordinary matter relativistic effects are not dominant but they are noticeable. In large atoms these effects severely change the innermost electrons and this has a noticeable effect on the overall electron density profile. Therefore, some version of relativistic mechanics is needed, which means, presumably, that we must know how to replace $p^2/2m$ by the Dirac operator.

The combination of relativistic mechanics plus the electromagnetic field (in addition to the Coulomb interaction) makes the stability problem difficult and uncertain. Major aspects of this problem have been worked out in the last few years (about 35) and that is the subject of this lecture.

3 Nonrelativistic Matter without the Magnetic Field

We work in the ‘Coulomb’ gauge for the electromagnetic field. Despite the assertion that quantum mechanics and quantum field theory are gauge invariant, it seems to be essential to use this gauge, even though its relativistic covariance is not as transparent as that of the Lorentz gauge. The reason is the following.

In the Coulomb gauge the electrostatic part of the interaction of matter with the electromagnetic field is put in ‘by hand’, so to speak. That is, it is represented by an ordinary potential αV_c , of the form (in energy units mc^2 and length units the Compton wavelength $\hbar/mc$)

$$V_c = - \sum_{i=1}^N \sum_{k=1}^K \frac{Z_k}{|\mathbf{x}_i - \mathbf{R}_k|} + \sum_{1 \leq i < j \leq N} \frac{1}{|\mathbf{x}_i - \mathbf{x}_j|} + \sum_{1 \leq k < l \leq K} \frac{Z_k Z_l}{|\mathbf{R}_k - \mathbf{R}_l|} . \quad (4)$$

The first sum is the interaction of the electrons (with dynamical coordinates $\mathbf{x}_i$) and fixed nuclei located at $\mathbf{R}_k$ of positive charge Z_k times the (negative) electron charge e . The second is the electron-electron repulsion and the third is the nucleus-nucleus repulsion. The nuclei are fixed because they are so massive relative to the electron that their motion is irrelevant. It could be included, however, but it would change nothing essential. Likewise there is no nuclear structure factor because if it were essential for stability then the size of atoms would be 10^{-13} cm instead of 10^{-8} cm, contrary to what is observed.

Although the nuclei are fixed the constant C in the stability of matter (2) is required to be independent of the $\mathbf{R}_k$ ’s. Likewise (1) requires that E_0 have a finite lower bound that is independent of the $\mathbf{R}_k$ ’s.

For simplicity of exposition we shall assume here that all the Z_k are identical, i.e., $Z_k = Z$.

The magnetic field, which will be introduced later, is described by a vector potential $\mathbf{A}(x)$ which is a dynamical variable in the Coulomb gauge. The

magnetic field is $\mathbf{B} = \text{curl}\mathbf{A}$.

There is a basic physical distinction between electric and magnetic forces which does not seem to be well known, but which motivates this choice of gauge. In electrostatics like charges repel while in magnetostatics like currents attract. A consequence of these facts is that the correct magnetostatic interaction energy can be obtained by minimizing the energy functional $\int B^2 + \int \mathbf{j} \cdot \mathbf{A}$ with respect to the vector field $\mathbf{A}$. The electrostatic energy, on the other hand, *cannot* be obtained by a minimization principle with respect to the field (e.g., minimizing $\int |\nabla\phi|^2 + \int \phi\rho$ with respect to ϕ).

The Coulomb gauge, which puts in the electrostatics correctly, by hand, and allows us to minimize the total energy with respect to the $\mathbf{A}$ field, is the gauge that gives us the correct physics and is consistent with the “quintessential quantum mechanical notion of a ground state energy” mentioned in Sect. 2.1. In any other gauge one would have to look for a critical point of a Hamiltonian rather than a true global minimum.

The type of Hamiltonian that we wish to consider in this section is

$$H_N = T_N + \alpha V_c . \quad (5)$$

Here, T is the kinetic energy of the N electrons and has the form

$$T_N = \sum_{i=1}^N T_i , \quad (6)$$

where T_i acts on the coordinate of the i^{th} electron. The nonrelativistic choice is $T = p^2$ with $\mathbf{p} = -i\nabla$ and $p^2 = -\Delta$.

3.1 Nonrelativistic Stability for Fermions

The problem of stability of the second kind for nonrelativistic quantum mechanics was recognized in the early days by a few physicists, e.g., Onsager, but not by many. It was not solved until 1967 in one of the most beautiful papers in mathematical physics by Dyson and Lenard [9].

They found that the Pauli principle, i.e., Fermi-Dirac statistics, is essential. Mathematically, this means that the Hilbert space is the subspace of antisymmetric functions, i.e., $\mathcal{H}^{\text{phys}} = \wedge^N L^2(\mathbb{R}^3; \mathbb{C}^2)$. This is how the Pauli principle is interpreted post-Schrödinger; Pauli invented his principle a year earlier, however!

Their value for C in (2) was rather high, about -10^{15} eV for $Z = 1$. The situation was improved later by Thirring and myself [27] to about -20 eV for $Z = 1$ by introducing an inequality that holds only for the kinetic energy of fermions (not bosons) in an arbitrary state Ψ .

$$\langle \Psi, T_N \Psi \rangle \geq (\text{const.}) \int_{\mathbb{R}^3} \varrho_\Psi(\mathbf{x})^{5/3} d^3\mathbf{x} , \quad (7)$$

where ϱ_Ψ is the one-body density in the (normalized) fermionic wave function Ψ (of space and spin) given by an integration over $(N - 1)$ coordinates and N spins as follows.

$$\varrho_\Psi(\mathbf{x}) = N \sum_{\sigma_1, \dots, \sigma_N} \int_{\mathbb{R}^{3(N-1)}} |\Psi(\mathbf{x}, \mathbf{x}_2, \dots, \mathbf{x}_N; \sigma_1, \dots, \sigma_N)|^2 d^3\mathbf{x}_2 \cdots d^3\mathbf{x}_N . \quad (8)$$

Inequality (7) allows one simply to reduce the quantum mechanical stability problem to the stability of Thomas-Fermi theory, which was worked out earlier by Simon and myself [26].

The older inequality of Sobolev,

$$\langle \Psi, T_N \Psi \rangle \geq (\text{const.}) \left(\int_{\mathbb{R}^3} \varrho_\Psi(\mathbf{x})^3 d^3\mathbf{x} \right)^{1/3} , \quad (9)$$

is not as useful as (7) for the many-body problem because its right side is proportional to N instead of $N^{5/3}$.

It is amazing that from the birth of quantum mechanics to 1967 none of the luminaries of physics had quantified the fact that electrostatics plus the uncertainty principle *do not suffice* for stability of the second kind, and thereby make thermodynamics possible (although they do suffice for the first kind). See Sect. 3.2. It was noted, however, that the Pauli principle was responsible for the large sizes of atoms and bulk matter (see, e.g., [8, 9]).

3.2 Nonrelativistic Instability for Bosons

What goes wrong if we have charged bosons instead of fermions? Stability of the first kind (1) holds in the nonrelativistic case, but (2) fails. If we assume the nuclei are infinitely massive, as before, and $N = KZ$ then $E_0 \sim -N^{5/3}$ [9, 18]. To remedy the situation we can let the nuclei have finite mass (e.g., the same mass as the negative particles). Then, as Dyson showed [8], $E_0 \leq -(\text{const.})N^{7/5}$. This calculation was highly non-trivial! Dyson had

to construct a variational function with pairing of the Bogolubov type in a rigorous fashion and this took several pages.

Thus, finite nuclear mass improves the situation, but not enough. The question whether $N^{7/5}$ is the correct power law remained open for many years. A lower bound of this type was needed and that was finally done in [5].

The results of this Section 3 can be summarized by saying that stability of the hydrogen atom is one thing but stability of many-body physics is something else !

4 Relativistic Kinematics (no magnetic field)

The next step is to try to get some idea of the effects of relativistic kinematics, which means replacing p^2 by $\sqrt{p^2 + 1}$ in non-quantum physics. The simplest way to do this is to substitute $\sqrt{p^2 + 1}$ for T in (6). The Dirac operator will be discussed later on, but for now this choice of T will suffice. Actually, it was Dirac's choice before he discovered his operator and it works well in some cases. For example, Chandrasekhar used it successfully, and accurately, to calculate the collapse of white dwarfs (and later, neutron stars).

Since we are interested only in stability, we may, and shall, substitute $|\mathbf{p}| = \sqrt{-\Delta}$ for T . The error thus introduced is bounded by a constant times N since $|\mathbf{p}| < \sqrt{p^2 + 1} < |\mathbf{p}| + 1$ (as an operator inequality). Our Hamiltonian is now $H_N = \sum_{i=1}^N |\mathbf{p}_i| + \alpha V_c$.

4.1 One-Electron Atom

The touchstone of quantum mechanics is the Hamiltonian for 'hydrogen' which is, in our case,

$$H = |\mathbf{p}| - Z\alpha/|\mathbf{x}| = \sqrt{-\Delta} - Z\alpha/|\mathbf{x}| . \quad (10)$$

It is well known (also to Dirac) that the analogous operator with $|\mathbf{p}|$ replaced by the Dirac operator ceases to make sense when $Z\alpha > 1$. Something similar happens for (10).

$$E_0 = \begin{cases} 0 & \text{if } Z\alpha \leq 2/\pi; \\ -\infty & \text{if } Z\alpha > 2/\pi . \end{cases} \quad (11)$$

The reason for this behavior is that both $|\mathbf{p}|$ and $|\mathbf{x}|^{-1}$ scale in the same way. Either the first term in (10) wins or the second does.

A result similar to (11) was obtained in [10] for the free Dirac operator $D(0)$ in place of $|\mathbf{p}|$, but with the wave function Ψ restricted to lie in the positive spectral subspace of $D(0)$. Here, the critical value is $\alpha Z \leq (4\pi)/(4 + \pi^2) > 2/\pi$.

The moral to be drawn from this is that relativistic kinematics plus quantum mechanics is a ‘critical’ theory (in the mathematical sense). This fact will plague any relativistic theory of electrons and the electromagnetic field – primitive or sophisticated.

4.2 Many Electrons and Nuclei

When there are many electrons is it true that the condition $Z\alpha \leq \text{const.}$ is the only one that has to be considered? The answer is no! One *also* needs the condition that α itself must be small, regardless of how small Z might be. This fact can be called a ‘discovery’ but actually it is an overdue realization of some basic physical ideas. It should have been realized shortly after Dirac’s theory in 1927, but it does not seem to have been noted until 1983 [7].

The underlying physical heuristics is the following. With α fixed, suppose $Z\alpha = 10^{-6} \ll 1$, so that an atom is stable, but suppose that we have 2×10^6 such nuclei. By bringing them together at a common point we will have a nucleus with $Z\alpha = 2$ and one electron suffices to cause collapse into it. Then (1) fails. What prevents this from happening, presumably, is the nucleus-nucleus repulsion energy which goes to $+\infty$ as the nuclei come together. But this repulsion energy is proportional to $(Z\alpha)^2/\alpha$ and, therefore, if we regard $Z\alpha$ as fixed we see that $1/\alpha$ must be large enough in order to prevent collapse.

Whether or not the reader believes this argument, the mathematical fact is that there is a fixed, finite number $\alpha_c \leq 2.72$ ([28]) so that when $\alpha > \alpha_c$ (1) fails for *every* positive Z and for every $N \geq 1$ (with or without the Pauli principle).

The open question was whether (2) holds for *all* N and K if $Z\alpha$ and α are both small enough. The breakthrough was due to Conlon [4] who proved (2), for fermions, if $Z = 1$ and $\alpha < 10^{-200}$. The situation was improved by Fefferman and de la Lave [12] to $Z = 1$ and $\alpha < 0.16$. Finally, the expected correct condition $Z\alpha \leq 2/\pi$ and $\alpha < 1/94$ was obtained in ([28]). (This paper contains a detailed history up to 1988.) The situation was further

improved in ([23]). The multi-particle version of the use of the free Dirac operator, as in Sect. 4.1, was treated in [16].

Finally, it has to be noted that charged bosons are *always* unstable of the first kind (not merely the second kind, as in the nonrelativistic case) for *every* choice of $Z > 0, \alpha > 0$. E.g., there is instability if $Z^{2/3}\alpha N^{1/3} > 36$ ([28]).

We are indeed fortunate that there are no stable, negatively charged bosons.

5 Interaction of Matter with Classical Magnetic Fields

The magnetic field $\mathbf{B}$ is defined by a vector potential $\mathbf{A}(\mathbf{x})$ and $\mathbf{B}(\mathbf{x}) = \text{curl } \mathbf{A}(\mathbf{x})$. In this section we take a first step (warmup exercise) by regarding $\mathbf{A}$ as classical, but indeterminate, and we introduce the classical field energy

$$H_f = \frac{1}{8\pi} \int_{\mathbb{R}^3} B(\mathbf{x})^2 dx . \quad (12)$$

The Hamiltonian is now

$$H_N(\mathbf{A}) = T_N(\mathbf{A}) + \alpha V_c + H_f , \quad (13)$$

in which the kinetic energy operator has the form (6) but depends on $\mathbf{A}$. We now define E_0 to be the infimum of $\langle \Psi, H_N(\mathbf{A})\Psi \rangle$ both with respect to Ψ and with respect to $\mathbf{A}$.

5.1 Nonrelativistic Matter with Magnetic Field

The simplest situation is merely ‘minimal coupling’ without spin, namely,

$$T(\mathbf{A}) = |\mathbf{p} + \sqrt{\alpha}\mathbf{A}(\mathbf{x})|^2 \quad (14)$$

This choice does not change any of our previous results qualitatively. The field energy is not needed for stability. On the one particle level, we have the ‘diamagnetic inequality’ $\langle \phi, |\mathbf{p} + \mathbf{A}(\mathbf{x})|^2 \phi \rangle \geq \langle |\phi|, p^2 |\phi| \rangle$. The same holds for $|\mathbf{p} + \mathbf{A}(\mathbf{x})|$ and $|\mathbf{p}|$. More importantly, inequality (7) for fermions continues to hold (with the same constant) with $T(\mathbf{A})$ in place of p^2 . (There

is an inequality similar to (7) for $|\mathbf{p}|$, with $5/3$ replaced by $4/3$, which also continues to hold with minimal substitution [6].)

The situation gets much more interesting if spin is included. This takes us a bit closer to the relativistic case. The kinetic energy operator is the Pauli-Fierz operator

$$T^P(\mathbf{A}) = |\mathbf{p} + \sqrt{\alpha} \mathbf{A}(\mathbf{x})|^2 + \sqrt{\alpha} \mathbf{B}(\mathbf{x}) \cdot \boldsymbol{\sigma} , \quad (15)$$

where $\boldsymbol{\sigma}$ is the vector of Pauli spin matrices.

5.1.1 One-Electron Atom

The stability problem with $T^P(\mathbf{A})$ is complicated, even for a one-electron atom. Without the field energy H_f the Hamiltonian is unbounded below. (For fixed $\mathbf{A}$ it is bounded but the energy tends to $-\infty$ like $-(\log B)^2$ for a homogeneous field [1].) The field energy saves the day, but the result is surprising [13] (recall that we must minimize the energy with respect to Ψ and $\mathbf{A}$):

$$|\mathbf{p} + \sqrt{\alpha} \mathbf{A}(\mathbf{x})|^2 + \sqrt{\alpha} \mathbf{B}(\mathbf{x}) \cdot \boldsymbol{\sigma} - Z\alpha/|\mathbf{x}| + H_f \quad (16)$$

is bounded below if and only if $Z\alpha^2 \leq C$, where C is some constant that can be bounded as $1 < C < 9\pi^2/8$.

The proof of instability [29] is difficult and requires the construction of a zero mode (soliton) for the Pauli operator, i.e., a finite energy magnetic field and a *square integrable* ψ such that

$$T^P(\mathbf{A})\psi = 0 . \quad (17)$$

The usual kinetic energy $|\mathbf{p} + \mathbf{A}(\mathbf{x})|^2$ has no such zero mode for any $\mathbf{A}$, even when 0 is the bottom of its spectrum.

The original magnetic field [29] that did the job in (17) is independently interesting, geometrically (many others have been found since then).

$$\mathbf{B}(x) = \frac{12}{(1+x^2)^3} [(1-x^2)\mathbf{w} + 2(\mathbf{w} \cdot \mathbf{x})\mathbf{x} + 2\mathbf{w} \wedge \mathbf{x}]$$

with $|\mathbf{w}| = 1$. The field lines of this magnetic field form a family of curves, which, when stereographically projected onto the 3-dimensional unit sphere, become the great circles in what topologists refer to as the Hopf fibration.

Thus, we begin to see that nonrelativistic matter with magnetic fields behaves like relativistic matter without fields – to some extent.

The moral of this story is that a magnetic field, which we might think of as possibly self-generated, can cause an electron to fall into the nucleus. The uncertainty principle cannot prevent this, not even for an atom!

5.1.2 Many Electrons and Many Nuclei

In analogy with the relativistic (no magnetic field) case, we can see that stability of the first kind fails if $Z\alpha^2$ or α are too large. The heuristic reasoning is the same and the proof is similar.

We can also hope that stability of the second kind holds if both $Z\alpha^2$ and α are small enough. The problem is complicated by the fact that it is the field energy H_f that will prevent collapse, but there there is only one field energy while there are $N \gg 1$ electrons.

The hope was finally realized, however. Fefferman [11] proved stability of the second kind for $H_N(\mathbf{A})$ with the Pauli-Fierz $T^P(\mathbf{A})$ for $Z = 1$ and “ α sufficiently small”. A few months later it was proved [24] for $Z\alpha^2 \leq 0.04$ and $\alpha \leq 0.06$. With $\alpha = 1/137$ this amounts to $Z \leq 1050$. This very large Z region of stability is comforting because it means that perturbation theory (in $\mathbf{A}$) can be reliably used for this particular problem.

Using the results in [24], Bugliaro, Fröhlich and Graf [2] proved stability of the same nonrelativistic Hamiltonian – but with an ultraviolet cutoff, quantized magnetic field whose field energy is described below. (Note: No cutoffs are needed for classical fields.)

There is also the very important work of Bach, Fröhlich, and Sigal [3] who showed that this nonrelativistic Hamiltonian with ultraviolet cutoff, quantized field *and* with sufficiently small values of the parameters has other properties that one expects. E.g., the excited states of atoms dissolve into resonances and only the ground state is stable. The infrared singularity notwithstanding, the ground state actually exists (the bottom of the spectrum is an eigenvalue); this was shown in [3] for small parameters and in [14] for all values of the parameters.

6 Relativity Plus Magnetic Fields

As a next step in our efforts to understand QED and the many-body problem we introduce relativity theory along with the classical magnetic field.

6.1 Relativity Plus Classical Magnetic Fields

Originally, Dirac and others thought of replacing $T^P(\mathbf{A})$ by $\sqrt{T^P(\mathbf{A}) + 1}$ but this was not successful mathematically and does not seem to conform to experiment. Consequently, we introduce the Dirac operator for T in (6), (13)

$$D(\mathbf{A}) = \boldsymbol{\alpha} \cdot \mathbf{p} + \sqrt{\alpha} \boldsymbol{\alpha} \cdot \mathbf{A}(\mathbf{x}) + \beta m , \quad (18)$$

where $\boldsymbol{\alpha}$ and β denote the 4×4 Dirac matrices and $\sqrt{\alpha}$ is the electron charge as before. (This notation of $\boldsymbol{\alpha}$ and α is not mine.) We take $m = 1$ in our units. The Hilbert space for N electrons is

$$\mathcal{H} = \wedge^N L^2(\mathbb{R}^3; \mathbb{C}^4) . \quad (19)$$

The well known problem with $D(\mathbf{A})$ is that it is unbounded below, and so we cannot hope to have stability of the first kind, even with $Z = 0$. Let us imitate QED (but without pair production or renormalization) by restricting the electron wave function to lie in the positive spectral subspace of a Dirac operator.

Which Dirac operator?

There are two natural operators in the problem. One is $D(0)$, the free Dirac operator. The other is $D(\mathbf{A})$ that is used in the Hamiltonian. In almost all formulations of QED the electron is defined by the positive spectral subspace of $D(0)$. Thus, we can define

$$\mathcal{H}^{\text{phys}} = P^+ \mathcal{H} = P_{i=1}^N \pi_i \mathcal{H} , \quad (20)$$

where $P^+ = P_{i=1}^N \pi_i$, and π_i is the projector of onto the positive spectral subspace of $D_i(0) = \boldsymbol{\alpha} \cdot \mathbf{p}_i + \beta m$, the free Dirac operator for the i^{th} electron. We then restrict the allowed wave functions in the variational principle to those Ψ satisfying

$$\Psi = P^+ \Psi \quad i.e., \quad \Psi \in \mathcal{H}^{\text{phys}} . \quad (21)$$

Another way to say this is that we replace the Hamiltonian (13) by $P^+ H_N P^+$ on $\mathcal{H}$ and look for the bottom of its spectrum.

It turns out that this prescription leads to disaster! While the use of $D(0)$ makes sense for an atom, it fails miserably for the many-fermion problem, as discovered in [25] and refined in [15]. The result is:

For all $\alpha > 0$ in (18) (with or without the Coulomb term αV_c) one can find N large enough so that $E_0 = -\infty$.

In other words, the term $\sqrt{\alpha} \boldsymbol{\alpha} \cdot \mathbf{A}$ in the Dirac operator can cause an instability that the field energy cannot prevent.

It turns out, however, that the situation is saved if one uses the positive spectral subspace of the Dirac operator $D(\mathbf{A})$ to define an electron. (This makes the concept of an electron $\mathbf{A}$ dependent, but when we make the vector potential into a dynamical quantity in the next section, this will be less peculiar since there will be no definite vector potential but only a fluctuating quantity.) The definition of the physical Hilbert space is as in (20) but with π_i being the projector onto the positive subspace of the full Dirac operator $D_i(\mathbf{A}) = \boldsymbol{\alpha} \cdot \mathbf{p}_i + \sqrt{\alpha} \boldsymbol{\alpha} \cdot \mathbf{A}(\mathbf{x}_i) + \beta m$. Note that these π_i projectors commute with each other and hence their product P^+ is a projector.

The result [25] for this model ((13) with the Dirac operator and the restriction to the positive spectral subspace of $D(\mathbf{A})$) is reminiscent of the situations we have encountered before:

If α and Z are small enough stability of the second kind holds for this model.

Typical stability values that are rigorously established [25] are $Z \leq 56$ with $\alpha = 1/137$ or $\alpha \leq 1/8.2$ with $Z = 1$.

6.2 Relativity Plus Quantized Magnetic Field

The obvious next step is to try to imitate the strategy of Sect. 6.1 but with the quantized $\mathbf{A}$ field. This was done recently in [22].

$$\mathbf{A}(\mathbf{x}) = \frac{1}{2\pi} \sum_{\lambda=1}^2 \int_{|\mathbf{k}| \leq \Lambda} \frac{\vec{\epsilon}_\lambda(\mathbf{k})}{\sqrt{|\mathbf{k}|}} \left[a_\lambda(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} + a_\lambda^*(\mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{x}} \right] d^3\mathbf{k}, \quad (22)$$

where Λ is the ultraviolet cutoff on the wave-numbers $|\mathbf{k}|$. The operators a_λ, a_λ^* satisfy the usual commutation relations

$$[a_\lambda(\mathbf{k}), a_\nu^*(\mathbf{q})] = \delta(\mathbf{k} - \mathbf{q}) \delta_{\lambda,\nu}, \quad [a_\lambda(\mathbf{k}), a_\nu(\mathbf{q})] = 0, \quad \text{etc} \quad (23)$$

and the vectors $\vec{\varepsilon}_\lambda(\mathbf{k})$ are two orthonormal polarization vectors perpendicular to $\mathbf{k}$ and to each other.

The field energy H_f is now given by a normal ordered version of (12)

$$H_f = \sum_{\lambda=1,2} \int_{\mathbb{R}^3} |\mathbf{k}| a_\lambda^*(\mathbf{k}) a_\lambda(\mathbf{k}) d^3\mathbf{k} \quad (24)$$

The Dirac operator is the same as before, (18). Note that $D_i(\mathbf{A})$ and $D_j(\mathbf{A})$ still commute with each other (since $\mathbf{A}(\mathbf{x})$ commutes with $\mathbf{A}(\mathbf{y})$). This is important because it allows us to imitate Sect. 6.1.

In analogy with (19) we define

$$\mathcal{H} = \wedge^N L^2(\mathbb{R}^3; \mathbb{C}^4) \otimes \mathcal{F} , \quad (25)$$

where $\mathcal{F}$ is the Fock space for the photon field. We can then define the *physical* Hilbert space as before

$$\mathcal{H}^{\text{phys}} = \Pi \mathcal{H} = \prod_{i=1}^N \pi_i \mathcal{H} , \quad (26)$$

where the projectors π_i project onto the positive spectral subspace of either $D_i(0)$ or $D_i(\mathbf{A})$.

Perhaps not surprisingly, the former case leads to catastrophe, as before. This is so, even with the ultraviolet cutoff, which we did not have in Sect. 6.1. Because of the cutoff the catastrophe is milder and involves instability of the second kind instead of the first kind. This result relies on a coherent state construction in [15].

The latter case (use of $D(\mathbf{A})$ to define an electron) leads to stability of the second kind if Z and α are not too large. Otherwise, there is instability of the second kind. The rigorous estimates are comparable to the ones in Sect. 6.1.

Clearly, many things have yet to be done to understand the stability of matter in the context of QED. Renormalization and pair production have to be included, for example.

The results of this section suggest, however, that a significant change in the Hilbert space structure of QED might be necessary. We see that it does not seem possible to keep to the current view that the Hilbert space is a simple tensor product of a space for the electrons and a Fock space for the photons. That leads to instability for many particles (or large charge, if the idea of ‘particle’ is unacceptable). The ‘bare’ electron is not really a good

physical concept and one must think of the electron as always accompanied by its electromagnetic field. Matter and the photon field are inextricably linked in the Hilbert space $\mathcal{H}^{\text{phys}}$.

The following tables [22] summarize the results of this and the previous sections

Electrons defined by projection onto the positive subspace of $D(0)$, the free Dirac operator

	Classical or quantized field without cutoff Λ $\alpha > 0$ but arbitrarily small.	Classical or quantized field with cutoff Λ $\alpha > 0$ but arbitrarily small.
Without Coulomb potential αV_c	Instability of the first kind	Instability of the second kind
With Coulomb potential αV_c	Instability of the first kind	Instability of the second kind

Electrons defined by projection onto the positive subspace of $D(A)$, the Dirac operator with field

	Classical field with or without cutoff Λ or quantized field with cutoff Λ
Without Coulomb potential αV_c	The Hamiltonian is positive
With Coulomb potential αV_c	Instability of the first kind when either α or $Z\alpha$ is too large
	Stability of the second kind when both α and $Z\alpha$ are small enough

References

- [1] J. Avron, I. Herbst, B. Simon: Schrödinger operators with magnetic fields III, Commun. Math. Phys. **79**, 529- 572 (1981).
- [2] L. Bugliaro, J. Fröhlich, G. M. Graf: Stability of quantum electrodynamics with nonrelativistic matter, Phys. Rev. Lett. **77**, 3494-3497 (1996).
- [3] V. Bach, J. Fröhlich, I. M. Sigal: Spectral analysis for systems of atoms and molecules coupled to the quantized radiation field. Comm. Math. Phys. **207** 249-290 (1999).
- [4] J. G. Conlon: The ground state of a classical gas, Commun. Math. Phys. **94**, 439-458 (1984).
- [5] J. G. Conlon, E. H. Lieb, H.-T. Yau: The $N^{7/5}$ law for charged bosons, Commun. Math. Phys. **116**, 417-448 (1988).
- [6] I. Daubechies: An uncertainty principle for fermions with generalized kinetic energy, Commun. Math. Phys. **90**, 511-520 (1983).
- [7] I. Daubechies, E. H. Lieb: One electron relativistic molecules with Coulomb interaction, Commun. Math. Phys. **90**, 497-510 (1983).
- [8] F. J. Dyson: Ground state energy of a finite system of charged particles, J. Math. Phys. **8**, 1538-1545 (1967).
- [9] F. J. Dyson, A. Lenard: Stability of matter I and II, J. Math. Phys. **8**, 423-434 (1967), **9**, 1538-1545 (1968).
- [10] W. D. Evans, P. P. Perry, H. Siedentop: The spectrum of relativistic one-electron atoms according to Bethe and Salpeter, Commun. Math. Phys. **178**, 733-746 (1996).
- [11] C. Fefferman: Stability of Coulomb systems in a magnetic field, Proc. Nat. Acad. Sci. USA, **92**, 5006-5007 (1995).
- [12] C. Fefferman, R. de la Llave: Relativistic stability of matter. I. Rev. Mat. Iberoamericana **2**, 119-213 (1986).

- [13] J. Fröhlich, E. H. Lieb, M. Loss: Stability of Coulomb systems with magnetic fields I. The one-electron atom, Commun. Math. Phys. **104**, 251-270 (1986). See also E. H. Lieb, M. Loss: Stability of Coulomb systems with magnetic Fields II. The many-electron atom and the one-electron molecule, Commun. Math. Phys. **104**, 271-282 (1986).
- [14] M. Griesemer, E. H. Lieb, M. Loss: Ground states in non-relativistic quantum electrodynamics, Invent. Math. **145**, 557-595 (2001).
- [15] M. Griesemer, C. Tix: Instability of pseudo-relativistic model of matter with self-generated magnetic field, J. Math. Phys. **40**, 1780-1791 (1999).
- [16] G. Hoever, H. Siedentop: The Brown-Ravenhall operator, Math. Phys. Electronic Jour. **5**, no. 6 (1999).
- [17] E. H. Lieb: The stability of matter, Rev. Mod. Phys. **48**, 553-569 (1976).
- [18] E. H. Lieb: The $N^{5/3}$ law for bosons, Phys. Lett. **70A**, 71-73 (1979).
- [19] E. H. Lieb: The stability of matter: From atoms to stars, Bull. Amer. Math. Soc. **22**, 1-49 (1990).
- [20] E. H. Lieb, J.L. Lebowitz: The existence of thermodynamics for real matter with Coulomb forces, Phys. Rev. Lett. **22**, 631-634 (1969).
- [21] E. H. Lieb, M. Loss: *Analysis*, American Mathematical Society (1997).
- [22] E. H. Lieb, M. Loss: Stability of a Model of Relativistic Quantum Electrodynamics, Commun. Math. Phys. **228**, 561-588 (2002). arXiv math-ph/0109002, mp_arc 01-315.
- [23] E. H. Lieb, M. Loss, H. Siedentop: Stability of relativistic matter via Thomas-Fermi theory, Helv. Phys. Acta **69**, 974-984 (1996).
- [24] E. H. Lieb, M. Loss, J. P. Solovej: Stability of matter in magnetic fields, Phys. Rev. Lett. **75**, 985-989 (1995).
- [25] E. H. Lieb, H. Siedentop, J. P. Solovej: Stability and instability of relativistic electrons in magnetic fields, J. Stat. Phys. **89**, 37-59 (1997). See also Stability of relativistic matter with magnetic fields, Phys. Rev. Lett. **79**, 1785-1788 (1997).

- [26] E. H. Lieb, B. Simon: Thomas-Fermi theory revisited, *Phys. Rev. Lett.* **31**, 681-683 (1973).
- [27] E. H. Lieb, W. Thirring: Bound for the kinetic energy of fermions which proves the stability of matter, *Phys. Rev. Lett.* **35**, 687-689 (1975). Errata **35**, 1116 (1975).
- [28] E. H. Lieb, H.-T. Yau: The stability and instability of relativistic matter, *Commun. Math. Phys.* **118**, 177-213 (1988). See also Many-body stability implies a bound on the fine structure constant, *Phys. Rev. Lett.* **61**, 1695-1697 (1988).
- [29] M. Loss, H.-T. Yau: Stability of Coulomb systems with magnetic fields III. Zero energy bound states of the Pauli operator, *Commun. Math. Phys.* **104**, 283-290 (1986).